

chemotherapy patients with folliculitis

This protocol is designed for chemotherapy patients who are experiencing folliculitis. The ingredients used in these products will gently reduce and prevent follicular impaction and inflammation.

Morning:

1. Cleanse the skin with **blemish control bar**. Rinse with warm water, and pat dry. (Consider using a cleansing tool like the Clarisonic® Skin Brush to lather and apply the cleansing bar.)
2. Apply one pump of **anti-redness serum** to reduce redness and calm irritation.
3. Apply two pumps of **hydrating serum** to maintain important moisture levels in the skin that minimize excessive dryness and discomfort.
4. For all over face and body care, apply **perfecting face & body hydrator SPF 30** for broad-spectrum UV protection and a hydrated, even skin tone.
5. For the lips, apply **peptide lip therapy** to reduce peeling, improve hydration and dramatically reduce lip lines.

Evening:

1. Cleanse the skin with **blemish control bar**. Rinse with warm water, and pat dry.
2. Apply a thin layer of **acne gel** (as needed) to areas of folliculitis breakouts.
3. Apply two pumps of **hydrating serum**.
4. Apply **ReBalance** for light hydration and additional calming and redness reduction. **silkcoat® balm** may be used in addition to or in place of **ReBalance** to relieve extreme dryness.
5. For the lips, apply **peptide lip therapy**.

As needed: après peel® soothing balm may be applied in seven-day intervals to reduce inflammation, irritation and histamine reactions.